CME QUESTIONS / STE SORULARI

1- Which paramaters of the below had been measured in the study of Yılmaz et al.

- a) Pelvic incidance
- **b)** Lumbar lordosis
- c) Thoracic kyphosis
- **d)** T-1 slop angle
- e) None of above
- 2- How many normal healthy patients had been evaluated in the study of Yılmaz et al?
 - **a)** 10
 - **b)** 20
 - **c)** 30
 - **d)** 40
 - **e)** 50
- 3- Which sentence of the below is true according to the second study of Yılmaz et al?
 - **a)** C0 inclination angles has been increased.
 - **b)** CO-C2 angle has been decreased.
 - c) T1 slope angle has been decreased.
 - d) Cervical lordosis has been unchanged.
 - **e)** Degenerative disc disease effect sagital plane parameters of the subaxial region

4- How many patients with the degenerative disc disease was evaluated in the second study of Yılmaz et al?

- **a)** 30
- **b)** 40
- **c)** 50
- **d)** 60
- **e)** 70
- 5- Which parameters below had been changed with the statistically significance according to the study of Er *et al*?
 - **a)** C0 inclination angle
 - **b)** CO-C2 angle
 - c) C7-T1 local sagittal contour
 - d) T1 slope angle
 - e) Cervical lordosis

6- How many male patients had been evaluated in the study of Er *et al*?

CME QUESTIONS / STE SORULARI

- **a)** 14
- **b)** 24
- **c)** 34
- **d)** 44
- **e)** 54

7- How many complex operation had been performed according to the study of Gerilmez *et al*?

- **a)** 17
- **b)** 37
- **c)** 37
- **d)** 47
- **e)** 57

8- Which sentence below was true according to the study of Gerilmez *et al*?

- a) It has been observed that surgical instruments may not be contaminated with airborne particles.
- **b)** Great attention must be given to appropriate behavior regarding contamination in the operating room,
- c) Activity must be kept to a minimum,
- d) Doors must be controlled.
- e) All of the sentence above
- 9- How many cases had been evaluated for the sexual function in the study of Onen *et al*?
 - **a)** 50
 - **b)** 40
 - **c)** 30
 - **d)** 20
 - **e)** 10
- 10-Which percentages of the patients with the sexual disfunction due to herniation of cervical disc below has been calculated in the study of Onen *et al*?
 - **a)** 17.2 %
 - **b)** 34.5 %
 - **c)** 58.6 %
 - **d)** 65.6 %
 - **e)** 78.1 %

JTSS 27(3) issue CORRECT ANSWERS OF CME QUESTIONS:

1.	d	6.	c
2.	d	7.	e
3.	b	8.	a
4.	e	9.	d
5.	d	10.	c