CME QUESTIONS / STE SORULARI

- 1- Which subject of the below had been tested in the study of Sariyilmaz et al.
 - a) Guina pigs
 - **b)** Rabbits
 - c) Rats
 - d) Chickens
 - e) Dogs
- 2- Which drug had been used for evaluating pathogenesis of idiopathic scoliosis in the study of Sariyilmaz et al?
 - a) Steroids
 - b) Melatonin
 - c) Growth hormone
 - d) Botulinum toxin-A
 - e) Ergocalcipherol
- 3- Which experimental spinal cord injury area in the rats of the below had been made in the study of Cömert et al?
 - a) Upper cervical region
 - **b)** Lower cervical region
 - c) Thoracic region
 - d) Thoracolumbar region
 - e) L-1 level
- 4- How many rats with the spinal cord injury had been evaluated in the study of Cömert et al?
 - **a)** 30
 - **b)** 40
 - **c)** 50
 - **d)** 60
 - **e)** 70
- 5- Which Lenke type below had been evaluated with the idiopathic scoliosis treated with selective fusion according to the second study of Sariyilmaz *et al*?
 - a) Type-I
 - **b)** Type-II
 - c) Type-III
 - d) Type-IV
 - e) Type-V

- 6- Which corrective maneuver had been evaluated for correction of the lumbar lordosis in the second study of Sariyilmaz *et al*?
 - a) Translation
 - b) Distraction
 - c) Compression
 - d) Convex rod derotation
 - e) None of all
- 7- How many balloon kyphoplasty had been performed according to the study of Demircay et al?
 - **a)** 23
 - **b)** 33
 - **c)** 43
 - **d)** 53
 - **e)** 63
- 8- Which sentence below was true according to the study of Demircay *et al*?
 - a) ODI scores is not changed improved at the final follow-up.
 - **b)** There were 15 female patients and 48 male patients.
 - c) Mean bone cement (PMMA) volume was 4.6 ml.
 - d) This study included 43 patients
 - e) All of the sentence above
- 9- How many cases had been evaluated in the study of Akyol et al?
 - **a)** 47
 - **b)** 37
 - **c)** 35
 - **d)** 27
 - **e)** 17
- 10- Which sentence below was true according to the study of Akyol *et al*?
 - a) Myelopathy symptoms were seen in 39 patients.
 - b) Radiculopathy symptoms were found in 18 patients.
 - c) ODL is an effective and safe way to reduce symptoms of degenerative cervical myelopathy due to OPLL
 - d) Preoperative mean mJOA scores were 1.4
 - e) None of all

JTSS 27(4) issue CORRECT ANSWERS OF CME QUESTIONS:

- 1. a 6. d
- 2. b 7. e
- 3. e 8. a
- 4. a 9. e
- 5. c 10.d