

- 1- In the study by Güçlü et al., which of the following is the vertebra for which a trauma model was constructed using the finite element method?
 - a) C1
 - b) C2
 - c) C3
 - d) C4
 - e) C5
- 2- Which of the following is incorrect according to the data obtained in the study by Oltulu et al.?
 - a) 30 patients were included in the study.
 - b) Two evaluators worked on the study.
 - c) A similar correlation was found between the manual measurement values and the computerized measurement values.
 - d) Sagittal parameter measurement with a computer was found to be more reliable than manual measurement.
 - e) A high consistency was found in the computerized measurements.
- 3- At which of the levels were the disc heights measured in the study by Köken et al.?
 - a) C1-2, C2-3, and C3-4
 - b) C3-4, T1-2, and L1-2
 - c) T11-12, T12-L1, and L1-2
 - d) C4-5, C5-6, and C6-7
 - e) L1-2, L2-3, and L4-5

- 4- What was the main benefit of the measurement of the disc heights in the study by Köken et al.?
 - a) Predetermination of disc degeneration
 - b) Evaluation of growth potential
 - c) Accurate determination of the movement borders
 - d) Application of the correct size of intravertebral cage and prosthesis
 - e) Determination of the severity of spinal stenosis
- 5- For how many patients was the atlantodental distance measured in the study by Özdoğan et al.?
 - a) 48
 - b) 60
 - c) 74
 - d) 88
 - e) 100
- 6- When the atlantodental distance is over which of the following values can pathological dissociation be considered, according to the results obtained in the study by Özdoğan et al.?
 - a) 10
 - b) 8
 - c) 6
 - d) 4
 - e) 2

- 7- Which of the following lengths should be preferred for transpedicular screws used in the thoracolumbar region in females, according to the study by Köken et al.?
 - a) 2-7 mm
 - b) 3-8 mm
 - c) 5-8 mm
 - d) 4-12 mm
 - e) 6-11 mm
- 8- How many patients received percutaneous endoscopic lumbar discectomy in the case report by Uğraş et al.?
 - a) 13
 - b) 23
 - c) 33
 - d) 43
 - e) 53
- 9- According to data obtained in the study by Uğraş et al., what percentage of patients recovered fully?
 - a) 40
 - b) 60
 - c) 70
 - d) 80
 - e) 100

- 10- At which vertebral levels were hemangiomas detected in the case report by Korkmaz et al.?
 - a) T4 and L4
 - b) T1 and L1
 - c) T6 and L2
 - d) T10 and L4
 - e) T12 and L2

JTSS 25(2) Number CORRECT ANSWERS OF CME QUESTIONS:

1. d	6. c
2. e	7. c
3. d	8. d
4. a	9. d
5.b	10.b